Ideal Hemoglobin Targets In Renal Anemia is around 11.5 g/dL.¹

Erythropoiesis-Stimulating Agents are Initiated in Dialysis Patient With Hemoglobin 9.0-10.0 g/dL.¹



Darbepoetin alfa shows the Similar efficacy as Epoetin alfa, at a reduced dose frequency²



References:

1. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Int Suppl.* 2012;2(4):299-310. **2.** Chen N, Xing C, Niu J, *et al.* Darbepoetin alfa injection versus epoetin alfa injection for treating anemia of Chinese hemodialysis patients with chronic kidney failure: A randomized, open-label, parallel-group, non-inferiority phase III trial. *Chronic Dis Transl Med.* 2022;8(1):59–70.

Darbepoetin alfa 25/40/60µg

LMRC Code: GGI-CO-A1-AQS-DCVN-BANNERS-K23-0695

For the use of a Registered Medical Practitioner, Hospital or Laboratory only.

All content including: text, images, audio, or other formats were created for medical educational purposes only. Offerings for continuing education are clearly identified and the appropriate target audience is identified. This communication is restricted for the use of HCPs and must not be shared in public domain. The opinions expressed in this video are solely the views of the speaker. The presentation may contain off label usage of drug(s) aimed at educational purpose only. Dr. Reddy's Laboratories Ltd including but not limited its Affiliates strictly does not recommend any off label usage of its marketing products. Prior to treatment kindly refer to the latest prescribing information approved by the local authorities. No part of this content may be used, reproduced, transmitted or stored in any form without the written permission of Dr. Reddy's.